

Protecting and improving the nation's health

Lincoln

District



This profile was published on 4th July 2017

Health Profile 2017

Health in summary

The health of people in Lincoln is varied compared with the England average. Lincoln is one of the 20% most deprived districts/unitary authorities in England and about 23% (3,900) of children live in low income families. Life expectancy for both men and women is lower than the England average.

Health inequalities

Life expectancy is 8.6 years lower for men and 7.5 years lower for women in the most deprived areas of Lincoln than in the least deprived areas.

Child health


In Year 6, 22.8% (204) of children are classified as obese, worse than the average for England. The rate of alcohol-specific hospital stays among those under 18 is 19*, better than the average for England. This represents 3 stays per year. Levels of GCSE attainment are worse than the England average.

Adult health

The rate of alcohol-related harm hospital stays is 630*. This represents 546 stays per year. The rate of self-harm hospital stays is 237*, worse than the average for England. This represents 254 stays per year. The rate of TB is better than average.

Local priorities

Priorities in Lincoln include reducing alcohol misuse, smoking and obesity. For more information see http://www.research-lincs.org.uk and http://www.lincolnshire.gov.uk



Contains National Statistics data © Crown copyright and database right 2017 Contains OS data © Crown copyright and database right 2017


This profile gives a picture of people's health in Lincoln. It is designed to help local government and health services understand their community's needs, so that they can work together to improve people's health and reduce health inequalities.

Visit <u>www.healthprofiles.info</u> for more profiles, more information and interactive maps and tools.



^{*} rate per 100,000 population

Population: summary characteristics



| | Males | Females | Persons | |
|---|-------|---------|---------|--|
| Lincoln (population in thousar | | | | |
| Population (2015): | 48 | 49 | 97 | |
| Projected population (2020): | 49 | 50 | 99 | |
| % people from an ethnic minority group: | 4.1% | * | 3.5% | |
| Dependency ratio (d | 48.3% | | | |

| Population (2015): | 27,029 | 27,757 | 54,786 |
|---|--------|--------|--------|
| Projected population (2020): | 28,157 | 28,706 | 56,862 |
| % people from an ethnic minority group: | 13.1% | 13.4% | 13.2% |
| Dependency ratio (de | 60.7% | | |

^{* -} value suppressed due to small numbers


The age profile and table present demographic information for the residents of the area and England. They include a 2014-based population projection (to 2020), the percentage of people from an ethnic minority group (Annual Population Survey, October 2014 to September 2015) and the dependency ratio.

The dependency ratio estimates the number of dependants in an area by comparing the number of people considered less likely to be working (children aged under 16 and those of state pension age or above) with the working age population. A high ratio suggests the area might want to commission a greater level of services for older or younger people than those areas with a low ratio.


- Lincoln 2015 (Male)
- England 2015
- Lincoln 2015 (Female)
- Lincoln 2020 estimate

Deprivation: a national view

The map shows differences in deprivation in this area based on national comparisons, using national quintiles (fifths) of the Index of Multiple Deprivation 2015 (IMD 2015), shown by lower super output area. The darkest coloured areas are some of the most deprived neighbourhoods in England.




This chart shows the percentage of the population who live in areas at each level of deprivation.



Least deprived quintile

Life expectancy: inequalities in this local authority


The charts show life expectancy for men and women in this local authority for 2013-15. The local authority is divided into local deciles (tenths) by deprivation (IMD 2015), from the most deprived decile on the left of the chart to the least deprived decile on the right. The steepness of the slope represents the inequality in life expectancy that is related to deprivation in this local area. If there was no inequality in life expectancy the line would be horizontal.



deprived, the value could not be calculated as the number of cases is too small.


Local average

England average



Health inequalities: changes over time

These charts provide a comparison of the changes in death rates in people under 75 (early deaths) between this area and England. Early deaths from all causes also show the differences between the most and least deprived local quintile in this area. Data from 2010-12 onwards have been revised to use IMD 2015 to define local deprivation quintiles (fifths), all prior time points use IMD 2010. In doing this, areas are grouped into deprivation quintiles using the Index of Multiple Deprivation which most closely aligns with time period of the data. This provides a more accurate way of discriminating changes between similarly deprived areas over time.



Local most deprived

Local inequality

Local least deprived

Health summary for Lincoln

The chart below shows how the health of people in this area compares with the rest of England. This area's result for each indicator is shown as a circle. The average rate for England is shown by the black line, which is always at the centre of the chart. The range of results for all local areas in England is shown as a grey bar. A red circle means that this area is significantly worse than England for that indicator; however, a green circle may still indicate an important public health problem.

| Signi | ficantly worse than England average | | England | • | al average | e [€] | England average | England |
|--------------------------------------|--|------------------------|----------------|---------------------|----------------------|----------------|---------------------|-------------|
| O Not s | ignificantly different from England average | | worst | | • | 25th | 75th | best |
| Signi | ficantly better than England average | | | | | centile | percentile | |
| O Not c | ompared | Period | Local | Local | F== | F== | | Гъя |
| Domain | Indicator | Репоа | Local count | Local value | Eng value | Eng worst | England range | Eng best |
| young Our communities | 1 Deprivation score (IMD 2015) | 2015 | n/a | 28.9 | 21.8 | 42.0 | 0 | 5.0 |
| | 2 Children in low income families (under 16s) | 2014 | 3,925 | 23.3 | 20.1 | 39.2 | | 6.6 |
| | 3 Statutory homelessness | 2015/16 | 11 | 0.3 | 0.9 | | | |
| | 4 GCSEs achieved | 2015/16 | 384 | 45.8 | 57.8 | 44.8 | • • | 78.7 |
| | 5 Violent crime (violence offences) | 2015/16 | 1,766 | 18.4 | 17.2 | 36.7 | | 4.5 |
| | 6 Long term unemployment | 2016 | 463 | 7.0 ^ ²⁰ | 3.7 ^ ²⁰ | 13.8 | • • | 0.4 |
| | 7 Smoking status at time of delivery | 2015/16 | x ¹ | x ¹ | 10.6 \$ ¹ | 26.0 | ♦ | 1.8 |
| | 8 Breastfeeding initiation | 2014/15 | 760 | x ¹ | 74.3 | 47.2 | | 92.9 |
| and s he | 9 Obese children (Year 6) | 2015/16 | 204 | 22.8 | 19.8 | 28.5 | | 9.4 |
| Children's and young people's health | 10 Admission episodes for alcohol-specific conditions (under 18s)† | 2013/14 - 15/16 | 10 | 18.5 | 37.4 | 121.3 | ♦ ○ | 10.5 |
| | 11 Under 18 conceptions | 2015 | 34 | 24.1 | 20.8 | 43.8 | ○ | 5.4 |
| e ud | 12 Smoking prevalence in adults | 2016 | n/a | 21.0 | 15.5 | 25.7 | | 4.9 |
| Adults' health and lifestyle | 13 Percentage of physically active adults | 2015 | n/a | 54.0 | 57.0 | 44.8 | • | 69.8 |
| hea A | 14 Excess weight in adults | 2013 - 15 | n/a | 66.1 | 64.8 | 76.2 | | 46.5 |
| Disease and poor health | 15 Cancer diagnosed at early stage | 2015 | 146 | 45.1 | 52.4 | 39.0 | 0 | 63.1 |
| | 16 Hospital stays for self-harm† | 2015/16 | 254 | 237.1 | 196.5 | 635.3 | | 55.7 |
| oor h | 17 Hospital stays for alcohol-related harm† | 2015/16 | 546 | 630.0 | 647 | 1,163 | \(\bigcirc\) | 374 |
| од ра | 18 Recorded diabetes | 2014/15 | 4,938 | 6.4 | 6.4 | 9.2 | ♦ ♦ | 3.3 |
| se a | 19 Incidence of TB | 2013 - 15 | 12 | 4.2 | 12.0 | 85.6 | KO | 0.0 |
| isea | 20 New sexually transmitted infections (STI) | 2016 | 492 | 731.4 | 795 | 3,288 | \bigcirc | 223 |
| | 21 Hip fractures in people aged 65 and over† | 2015/16 | 90 | 592.6 | 589 | 820 | (| 312 |
| Life expectancy and causes of death | 22 Life expectancy at birth (Male) | 2013 - 15 | n/a | 77.6 | 79.5 | 74.3 | • • | 83.4 |
| | 23 Life expectancy at birth (Female) | 2013 - 15 | n/a | 81.9 | 83.1 | 79.4 | | 86.7 |
| | 24 Infant mortality | 2013 - 15 | 8 | 2.1 | 3.9 | 8.2 | ♦ | 0.8 |
| | 25 Killed and seriously injured on roads | 2013 - 15 | 100 | 34.6 | 38.5 | 103.7 | O | 10.4 |
| | 26 Suicide rate | 2013 - 15 | 35 | 13.7 | 10.1 | 17.4 | | 5.6 |
| | 27 Smoking related deaths | 2013 - 15 | n/a | n/a | 283.5 | | | |
| | 28 Under 75 mortality rate: cardiovascular | 2013 - 15 | 213 | 105.6 | 74.6 | 137.6 | | 43.1 |
| ехре | 29 Under 75 mortality rate: cancer | 2013 - 15 | 328 | 161.0 | 138.8 | 194.8 | | 98.6 |
| Life | 30 Excess winter deaths | Aug 2012 - Jul 2015 | 274 | 36.0 | 19.6 | 36.0 | | 6.9 |

Indicator notes

1 Index of Multiple Deprivation (IMD) 2015 2 % children (under 16) in low income families 3 Eligible homeless people not in priority need, crude rate per 1,000 households 4 5 A*-C including English & Maths, % pupils at end of key stage 4 resident in local authority 5 Recorded violence against the person crimes, crude rate per 1,000 population 6 Crude rate per 1,000 population aged 16-64 7 % of women who smoke at time of delivery 8 % of all mothers who breastfeed their babies in the first 48hrs after delivery 9 % school children in Year 6 (age 10-11) 10 Persons under 18 admitted to hospital due to alcohol-specific conditions, crude rate per 100,000 population 11 Under-18 conception rate per 1,000 females aged 15 to 17 (crude rate) 12 Current smokers (aged 18 and over), Annual Population Survey 13 % adults (aged 16 and over) achieving at least 150 mins physical activity per week, Active People Survey 14 % adults (aged 16 and over) classified as overweight or obese, Active People Survey 15 Experimental statistics - % of cancers diagnosed at stage 1 or 2 16 Directly age sex standardised rate per 100,000 population 17 Admissions involving an alcohol-related primary diagnosis or an alcohol-related external cause (narrow definition), directly age standardised rate per 100,000 population 18 % people (aged 17 and over) on GP registers with a recorded diagnosis of diabetes 19 Crude rate per 100,000 population 20 All new diagnoses (excluding chlamydia under age 25), crude rate per 100,000 population aged 15 to 64 21 Directly age-sex standardised rate of emergency admissions, per 100,000 population aged 65 and over 22, 23 The average number of years a person would expect to live based on contemporary mortality rates 24 Rate of deaths in infants aged under 1 year per 1,000 live births 25 Rate per 100,000 population 26 Directly age standardised rate per 100,000 population aged 35 and over 28 Directly age standardised rate per 100,000 population aged under 75 30 Ratio of excess winter deaths (observed winter deaths minus expe

† Indicator has had methodological changes so is not directly comparable with previously released values. € "Regional" refers to the former government regions.

x²0 Value based on an average of monthly counts x¹ Value not published for data quality reasons \$1 There is a data quality issue with this value

If 25% or more of areas have no data then the England range is not displayed.

Please send any enquiries to healthprofiles@phe.gov.uk

You may re-use this information (not including logos) free of charge in any format or medium, under the terms of the Open Government Licence. To view this licence, visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3/